

4-th Asian–Pacific Mathematical Olympiad 1992

1. A triangle with sides a, b, c is given. If its semiperimeter is denoted by s , we construct a triangle with sides $s - a$, $s - b$, and $s - c$. This process is repeated until a triangle can no longer be constructed with the side lengths given. For which original triangles can this process be repeated indefinitely?
2. In a circle C with center O and radius r , let C_1, C_2 be two circles with centers O_1, O_2 and radii r_1, r_2 respectively, so that each circle C_i is internally tangent to C at A_i and so that C_1, C_2 are externally tangent to each other at A . Prove that the three lines OA, O_1A_2 , and O_2A_1 are concurrent.
3. Let $n > 3$ be an integer. We select three numbers from the set $\{1, 2, \dots, n\}$. Using each of these three numbers only once and using addition, multiplication, and parenthesis, let us form all possible combinations.
 - (a) Show that if all three selected numbers are greater than $n/2$, then the values of these combinations are all distinct.
 - (b) Let $p \leq \sqrt{n}$ be a prime number. Show that the number of ways of choosing three numbers so that the smallest one is p and the values of the combinations are not all distinct is precisely the number of positive divisors of $p - 1$.
4. Determine all pairs (h, s) of positive integers with the following property:
If one draws h horizontal lines and another s lines which satisfy
 - (i) they are not horizontal,
 - (ii) no two of them are parallel, and
 - (iii) no three of the $h + s$ lines are concurrent,then the number of regions formed by these $h + s$ lines is 1992.
5. Find a sequence of maximal length consisting of non-zero integers in which the sum of any seven consecutive terms is positive and that of any eleven consecutive terms is negative.