

1st BALKAN STUDENT MATHEMATICAL COMPETITION

1. Matematičko natjecanje učenika Balkana

November 2008.

1st grade

Solutions

Problem 1. Find all nonnegative real numbers a and b for which

$$[a^3 + b^3 + (a + b)^3]^2 = 100a^3b^3.$$

(Adrian Satja Kurdija)

Solution. Let us notice that

$$\begin{aligned} a^3 + b^3 &\geq 2 \cdot a^{\frac{3}{2}} \cdot b^{\frac{3}{2}} \iff \left(a^{\frac{3}{2}} - b^{\frac{3}{2}}\right)^2 \geq 0, \\ (a + b)^3 &\geq 8 \cdot a^{\frac{3}{2}} \cdot b^{\frac{3}{2}} \iff a + b \geq 2 \cdot a^{\frac{1}{2}} \cdot b^{\frac{1}{2}} \iff \left(a^{\frac{1}{2}} - b^{\frac{1}{2}}\right)^2 \geq 0. \end{aligned}$$

(1 + 1 point)

Also, let us notice that it is possible to achieve equality in these two inequalities if and only if $a = b$. By adding these two inequalities, we gather that

$$[a^3 + b^3 + (a + b)^3]^2 \geq \left(2 \cdot a^{\frac{3}{2}} \cdot b^{\frac{3}{2}} + 8 \cdot a^{\frac{3}{2}} \cdot b^{\frac{3}{2}}\right)^2 = 100a^3b^3.$$

(3 points)

So, we achieve equality in the problem if and only if $a = b$ (it is necessary, and easy, to check that equality is achieved if $a = b$). Therefore, the only solutions to this equation are pairs $(a, b) = (c, c)$, where c can be any nonnegative real number.

(5 points)



Problem 2. In the forest, there are n trees lined up in a row, with n being a natural number greater than 1. To each tree there are assigned two numbers in the following way: to the tree in the i -th place ($i = 1, 2, \dots, n$), numbers assigned are a_i i b_i . b_i is equal to either 1 or -1 , while a_i can be any integer **not equal to 0**.

Horatio the Monkey stands on the k -th tree ($k \in \{1, 2, \dots, n\}$). He wishes to visit all n trees, but in such a manner that he visits each tree **exactly once**, jumping according to the following rule: From m -th ($m \in \{1, 2, \dots, n\}$) tree, he jumps $a_1b_1 + a_2b_2 + \dots + a_mb_m$ places. Jumping r places means jumping from the m -th tree to the $(m + r)$ -th tree (r is not necessarily positive). If $m + r < 1$ or $m + r > n$, Horatio will stay on the m -th tree and will jump no more.

Is it possible for Horatio, for any choice of numbers b_i , to choose numbers a_i so that he fulfills his goal? (Before jumping he assigns a certain number to each tree and than starts to jump, meaning he can not change numbers a_i during his jumping.)

(Melkior Ornik)

Solution. Firstly, let us notice that without loss of generality, we can assume $b_i = 1$ for every $i = 1, 2, \dots, n$. We are only interested in products $a_i \cdot b_i$, so, if some $b_i = -1$, we simply change the

sign of a_i . Further, for $n = 2$, the answer to the question in the problem is obviously positive: we let $a_1 = 1$ and $a_2 = -2$. (1 point)

Now we will show that the answer is positive for any $n \in \mathbb{N}$, $n > 2$. Let $c_i = a_1 + a_2 + \dots + a_i$. Let's look at three cases.

1° $k = 1$, so, Horatio "starts" from the first tree. We have the following two cases:

- $n = 2t + 1$, t being a natural number. In that case, for every $j \in \{1, 2, \dots, t\}$ we let $c_j = n - 2j + 1$ and for every $j \in \{t + 2, t + 3, \dots, 2t + 1\}$ we let $c_j = n - 2j + 2$. We will let $c_{t+1} = 0$. Now we can see that Horatio jumps from j -th tree to $(n - j + 1)$ -st tree when $j \in \{1, 2, \dots, t\}$ or to $(n - j + 2)$ -nd tree when $j \in \{t + 2, t + 3, \dots, 2t + 1\}$. So, he will jump from the first tree to the last one, from the last one to the second and so forth until, at the end, he jumps from the tree immediately right to the middle of the middle tree (the middle tree is $(t + 1)$ -st tree). We have, therefore, shown that Horatio will succeed in his goal. What we still need to show is that $a_i \neq 0$ for all $i \in \{2, 3, \dots, t\}$. (For $a_1 = c_1 = n - 1 \neq 0$ it is directly true.) Furthermore, for all $i \in \{2, 3, \dots, t\}$, $a_i = c_i - c_{i-1}$. However, it is obvious that there does not exist i such that $c_i = c_{i-1}$. With this, we have shown that Horatio can fulfill his goal in this case. (3 points)

- $n = 2t$, where t is a natural number greater than 1. Then, for every $j \in \{1, 2, \dots, t\}$ we let $c_j = n - 2j + 1$ and for every $j \in \{t + 1, t + 2, \dots, 2t\}$ we let $c_j = n - 2j + 2$. Now, we can analogously draw the same conclusions as in the previous case. (1 point)

2° $k = n$, which means Horatio "starts" from the last tree. This situation is analogous to the situation where $k = 1$ – Horatio only moves in the opposite direction all the time. So, in this case too Horatio can succeed in his goal. (1 point)

3° $1 < k < n$, which means Horatio "starts" from some tree which is neither the first nor the last one. Let us observe trees starting with k -th and ending with n -th tree. We can imagine that k is the first tree and then, according to the first case, we can see that he can visit them all in a permitted way. The tree that Horatio will visit last is easily found (in the first case we can see how). We can easily obtain a_i for that tree such that, from it, Horatio jumps to the $(k - 1)$ -th tree. Now, let us observe trees starting with the first one and ending with $k - 1$ -th one. According to the second case, Horatio can visit them all in a permitted way (if we look at $(k - 1)$ if it were the last tree). (3 points)


Finally, we have shown that Horatio can always succeed in his goal. (1 point) ■

Problem 3. An acute-angled triangle ABC is given. Bisector of angle $\angle BAC$ intersects the side \overline{BC} in point K . Point L is the midpoint of segment \overline{AK} and point N is the intersection of altitude from vertex A and side \overline{BC} . On side \overline{AB} , point P is such that $\angle KLP = \angle BAC$. Circumcircle of triangle ANP intersects side \overline{AC} in point Q . Prove that $|AP| = |AQ|$. (Adrian Satja Kurdija)

Solution. Let k be a circle with center in L and diameter \overline{AK} . By Thales' theorem on a right triangle AKN , point N is on the circle k . (1 point)

Let $\angle BAC = \angle KLP = 2\alpha$. Then, $\angle PAL = \alpha$ (angle bisector). Angle $\angle KLP$ is an exterior angle of triangle APL , so we can easily see that $\angle APL = \angle KLP - \angle PAL = 2\alpha - \alpha = \alpha$. So, $|LA| = |LP|$, and, as \overline{LA} is a radius of k , we can conclude that point P also lies on the circle k . Therefore, $\angle APK = 90^\circ$. (3 points)

We have shown that points A , N and P lie on k , and, because of that, k is the circumcircle of triangle ANP . So, point Q lies on circle k . \overline{AK} is a diameter of k so it follows that angle $\angle AQK = 90^\circ$. (2 points)



Now we can easily see that triangles APK and AQK are congruent (\overline{AK} is their common side, they're both right and they both have one acute angle of the same size). We conclude that $|AP| = |AQ|$. (4 points) ■

Problem 4. Find all natural numbers x, y i z and a prime number p , which is of the form $4k + 1$ ($k \in \mathbb{N}$), for which

$$4^x + p^y = 15^z.$$

(Melkior Ornik)

Solution. Let us denote the left side of this equality with L and the right with D . Because of $p \equiv 1 \pmod{4}$, $L \equiv 1 \pmod{4}$ and, therefore, $D \equiv 1 \pmod{4}$ must be true, which obviously means $15^z \equiv 1 \pmod{4}$. Since $15 \equiv -1 \pmod{4}$, we conclude that z must be even, so $z = 2z_1$, where $z_1 \in \mathbb{N}$. (1 point)

Now we have

$$p^y = (15^{z_1} - 2^x)(15^{z_1} + 2^x),$$

and since p is prime, we can deduce that $15^{z_1} - 2^x = p^\alpha$ and $15^{z_1} + 2^x = p^\beta$, for α and β which are some integers for which $0 \leq \alpha < \beta \leq y$. (1 point)

Now, let us look at two cases.

1° $\alpha > 0$. Then, we have $p \mid p^\beta - p^\alpha$, and since $p^\beta - p^\alpha = 2^{x+1}$, we have $p \mid 2^{x+1}$, which means $p = 2$. However, that is impossible since $p \equiv 1 \pmod{4}$. (1 point)

2° $\alpha = 0$. Then, $15^{z_1} - 2^x = 1$. If $x = 1$, we have $15^{z_1} = 3$, which, obviously, does not give any solutions. So, we have $x \geq 2$ and $1 = 15^{z_1} - 2^x \equiv 15^{z_1} \pmod{4}$. Since $15 \equiv -1 \pmod{4}$, we conclude that z_1 is even. Let $z_1 = 2z_2$ for some natural number z_2 . (1 point)

Now, we have

$$(15^{z_2} - 1)(15^{z_2} + 1) = 2^x.$$

It is obvious that $15^{z_2} - 1 = 2^m$ and $15^{z_2} + 1 = 2^n$, where m and n are integers for which $0 \leq m < n \leq x$. By subtracting given equalities, we gather $2^n - 2^m = 2$. The only two powers of 2 which differ by 2 are 4 and 2, so $15^{z_2} = 3$. However, this, once again, obviously gives no solutions. (2 points)

Hence, we have shown that there are no solutions in natural numbers. (4 points) ■

1st BALKAN STUDENT MATHEMATICAL COMPETITION

1. Matematičko natjecanje učenika Balkana

November 2008.

2nd grade

Solutions

Problem 1. If x , y and z are positive real numbers for which $x + y + z = 1$, prove the inequality

$$\frac{1}{\sqrt{x+y}} + \frac{1}{\sqrt{y+z}} + \frac{1}{\sqrt{z+x}} \leq \frac{1}{\sqrt{2xyz}}.$$

(Adrian Satja Kurdija)

Solution. Multiplying the given inequality by $2\sqrt{xyz}$, we get an equivalent inequality

$$2\sqrt{\frac{xyz}{x+y}} + 2\sqrt{\frac{xyz}{y+z}} + 2\sqrt{\frac{xyz}{z+x}} \leq \sqrt{2}.$$

Let us notice that $\frac{xy}{x+y} \leq \frac{x+y}{4} \iff (x-y)^2 \geq 0$, so $\frac{xy}{x+y} \leq \frac{x+y}{4}$. Now,

$$2\sqrt{\frac{xyz}{x+y}} = 2\sqrt{\frac{xy}{x+y} \cdot z} \leq 2\sqrt{\frac{x+y}{4} \cdot z} = \sqrt{z(x+y)}.$$

(2 points)

Using Arithmetic Mean - Geometric Mean inequality (or using the fact that the square of a real number is always nonnegative) on numbers $z\sqrt{2}$ and $\frac{x+y}{\sqrt{2}}$, we get

$$\sqrt{z(x+y)} = \sqrt{z\sqrt{2} \cdot \frac{x+y}{\sqrt{2}}} \leq \frac{1}{2} \left(z\sqrt{2} + \frac{x+y}{\sqrt{2}} \right) = \frac{(x+y+2z)\sqrt{2}}{4}.$$

(3 points)

With this, we have shown that

$$2\sqrt{\frac{xyz}{x+y}} \leq \frac{(x+y+2z)\sqrt{2}}{4}.$$

Analogously, we show that

$$\begin{aligned} 2\sqrt{\frac{xyz}{y+z}} &\leq \frac{(y+z+2x)\sqrt{2}}{4}, \\ 2\sqrt{\frac{xyz}{z+x}} &\leq \frac{(z+x+2y)\sqrt{2}}{4}. \end{aligned}$$

By adding these three inequalities we get

$$2\sqrt{\frac{xyz}{x+y}} + 2\sqrt{\frac{xyz}{y+z}} + 2\sqrt{\frac{xyz}{z+x}} \leq \frac{(4x+4y+4z)\sqrt{2}}{4} = \sqrt{2}.$$

We have, hence, proven the inequality in question.

(5 points)



Problem 2. A natural number is written in each cell of 10×10 table. It is known that, no matter which 5 columns and 5 rows of this table we choose, the sum of numbers in their 25 intersection cells is even. Prove that all the numbers in the table are even.

Solution. Let us first prove the following lemma.

Lemma 1. If the sum of each 5 of the given 10 natural numbers is even, then all these numbers are even.

Proof. Let's assume the opposite. It is clear that not all numbers can be odd. Therefore, there has to be at least one odd and at least one even number. Then, if there are 5 or more odd numbers, by choosing 5 odd numbers we reach a contradiction. If there are less than 5 odd numbers, by choosing 4 even and one odd number, we also reach a contradiction. Hence, this lemma is proven. (2 points)



Let us observe any 5 columns of the given table. For every row of the table, let's compute the sum of numbers in cells which we get by intersecting the row with these 5 columns. That way we get 10 natural numbers (one for each row). The sum of any 5 of these numbers is even (this follows from the conditions of the problem). Now, using **Lemma 1**, we conclude that each of these 10 sums is even. (4 points)

By observing all possible choices of 5 columns of the given table, we get that the sum of each 5 numbers of every row is even. Again, using **Lemma 1**, we conclude that every number in each row is even. Therefore, all numbers in the table is even. (4 points)




Problem 3. Let M_n and N_n be points on sides \overline{CA} and \overline{CB} of triangle ABC , respectively, such that

$$|CM_n| = \frac{1}{n} |CA|, \quad |CN_n| = \frac{1}{n+1} |CB|, \quad \forall n \in \mathbb{N}.$$

Find the locus of points $M_iN_i \cap M_jN_j$, where i and j are different natural number.

Note. $M_iN_i \cap M_jN_j$ denotes the intersection of lines M_iN_i and M_jN_j .

Solution. We intend to show that all the mentioned lines intersect in one point. That will be point D such that quadrilateral $ABDC$ is a parallelogram. (1 point)



Let n be any natural number. We draw line p parallel to line AC such that B is on p . Let D be the intersection of p and M_nN_n . Let us notice that $\angle M_nCN_n = \angle ACB = \angle CBD = \angle N_nBD$ (alternate interior angles) and $\angle CN_nM_n = \angle BN_nD$ (vertical angles). So, we have shown that triangles CM_nN_n and BDN_n are similar (they have two equal angles). (2 points)

Now,

$$\frac{|BD|}{|CM_n|} = \frac{|BN_n|}{|CN_n|},$$

which means

$$\begin{aligned} |BD| &= \frac{|CM_n| \cdot |BN_n|}{|CN_n|} = \frac{\frac{1}{n} \cdot |CA| \cdot (|CB| - |CN_n|)}{|CN_n|} \\ &= \frac{\frac{1}{n} |CA| \cdot (|CB| - \frac{1}{n+1} |CB|)}{\frac{1}{n+1} |CB|} \\ &= \frac{\frac{1}{n} |CA| \cdot \frac{n}{n+1} |CB|}{\frac{1}{n+1} |CB|} \\ &= |CA|. \end{aligned}$$

With this, we have shown that $|BD| = |AC|$ and, since we know that $BD \parallel AC$, it follows that quadrilateral $ABDC$ is a parallelogram. (4 points)

Now we know that line $M_n N_n$ goes through point D such that quadrilateral $ABDC$ is a parallelogram for each natural number n . Finally, we conclude that the locus of $M_i N_i \cap M_j N_j$, where i and j are different natural numbers, is point D such that quadrilateral $ABDC$ is a parallelogram. (3 points) ■

Problem 4. a , b and c are natural numbers. It is known that $a^2 + b^2 + abc$ has no more than 2008 natural divisors and that it is divisible by $(c + 2)^{1004}$. Prove that a and b are not relatively prime.

(Adrian Satja Kurdija)

Solution. Let $A = a^2 + b^2 + abc$ and $p = c + 2$, having in mind that then $p \geq 3$. Let's assume that there exists a prime number q such that $q^2 \mid p$. Then, $q^{2008} \mid A$, and as q^{2008} itself has $2009 > 2008$ divisors, we reach a contradiction. So, there does not exist a prime number q such that $q^2 \mid p$. Further, let us assume that there exist different prime numbers r and s which divide p . Then, $r^{1004} s^{1004} \mid A$ and, obviously, $1005^2 > 2008$ and we, again, have a contradiction. Hence, there do not exist two different prime numbers which both divide p . With all of this, we have shown that p is prime. (2 points)

Now, $c = p - 2$, and, for that reason,

$$A = a^2 + b^2 + abc = a^2 + b^2 + ab(p - 2) = a^2 - 2ab + b^2 + abp = (a - b)^2 + abp.$$

Since $p \mid A$ and $p \mid abp$, it follows that $p \mid (a - b)^2$ and, since p is prime, it further follows that $p \mid a - b$. From this, we conclude finally that $p^2 \mid (a - b)^2$. (2 points)

Furthermore, $p^2 \mid A$ and $p^2 \mid (a - b)^2$, so $p^2 \mid abp$, which leads to $p \mid ab$, must also be true. (1 point)

Now we know that $p \mid a - b$, $p \mid ab$ and that p is prime. Since $p \mid ab$, this means that p divides at least one of the numbers a and b . (1 point)

Without loss of generality, we may assume that $p \mid a$. Then, from $p \mid a - b$, it directly follows that $p \mid b$. So, numbers a and b are not relatively prime. (4 points) ■

1st BALKAN STUDENT MATHEMATICAL COMPETITION

1. Matematičko natjecanje učenika Balkana

November 2008.

3rd and 4th grade

Solutions

Problem 1. Find all functions $f: \mathbb{R} \rightarrow \mathbb{R}$ such that for every two real numbers x and y ,

$$f(f(x) + xy) = f(x) \cdot f(y + 1).$$

(Marko Radovanović)

Solution. Obvious solutions are $f(x) \equiv 0, \forall x \in \mathbb{R}; f(x) \equiv 1, \forall x \in \mathbb{R}$ and $f(x) \equiv x, \forall x \in \mathbb{R}$.

(1 point)

Let $x := 0, y := t - 1$. Then, $f(f(0)) = f(0)f(t), \forall t \in \mathbb{R}$.

(1 point)

Let us consider two cases.

1° $f(0) \neq 0$. Then, for each $t \in \mathbb{R}$, it is true that $f(t) = \frac{f(f(0))}{f(0)} = k$.

(1 point)

By plugging $t = 0$, we reach $f(0)^2 = f(f(0))$ or $k^2 = k$. If $k = 0$, we get $f(0) = 0$, which gives a contradiction. So, $k = 1$ must hold. In that case, we reach this solution: $f(x) \equiv 1, \forall x \in \mathbb{R}$.

(2 points)

2° $f(0) = 0$. Now we shall consider following two cases:

- $f(x) = 0$ if and only if $x = 0$. Then, if we plug $y := -1$, we reach

$$f(f(x) - x) = f(x)f(0) = 0 \implies f(x) - x = 0 \implies f(x) = x.$$

So, in this case we reach this solution: $f(x) \equiv x, \forall x \in \mathbb{R}$.

(2 points)

- There exists a real number $x_0 \neq 0$ such that $f(x_0) = 0$. Then, let $x := x_0$ and $y := \frac{t}{x_0}$ for some real number t . Then,

$$f\left(f(x_0) + x_0 \cdot \frac{t}{x_0}\right) = f(x_0)f\left(\frac{t}{x_0} + 1\right) = 0 \implies f(t) = 0.$$

Finally, we reach the solution $f(x) \equiv 0, \forall x \in \mathbb{R}$.

(2 points)

We can see that functions mentioned in the beginning (and only those functions) satisfy the conditions of the problem.

(1 point)



Problem 2. Paralampius the Gnu stands on number 1 on number line. He wants to come to a natural number k by a sequence of consecutive jumps. Let us denote the number of ways on which Paralampius can come from number 1 to number k with $f(k)$ ($f: \mathbb{N} \rightarrow \mathbb{N}_0$). Specially, $f(1) = 0$. A way is a sequence of numbers (with order) which Paralampius has visited on his travel from number 1 to number k . Paralampius can, from number b , jump to number

- $2b$ (always),
- $3b$ (always),
- b^2 (if $\frac{b^4}{6k} \in \mathbb{N}$, where k is a natural number on which he wants to come to in the end).

Prove that, for every natural number n , there exists a natural number m_0 such that for every natural number $m > m_0$,

$$f(m) < 2^{\alpha_1 + \alpha_2 + \dots + \alpha_i - n},$$

where $m = p_1^{\alpha_1} \cdot p_2^{\alpha_2} \cdot \dots \cdot p_i^{\alpha_i}$ ($p_1 < p_2 < \dots < p_i$ are prime divisors of number m and $i, \alpha_1, \alpha_2, \dots, \alpha_i$ are natural numbers) is a prime factorization of natural number m . It is known that this factorization is unique for every natural number $m > 1$.

(Melkior Ornik, Ivan Krijan)

Solution. Let's notice that Paralampius can only reach numbers of the form $2^a \cdot 3^b$ ($a, b \in \mathbb{N}_0$). That is because neither one of the allowed jumps "introduces" a prime factor different than 2 or 3 (first jump "introduces" 2, second one "introduces" 3, while the last one doubles the number of each of already existing prime numbers). Therefore, for each natural number m divisible by a prime number $p > 3$, it holds that $f(m) = 0 < 2^{\alpha_1 + \alpha_2 + \dots + \alpha_i - n}$, $\forall n \in \mathbb{N}$. Specially, the same claim holds for $f(1)$. Let us continue by observing only numbers of the form $2^a \cdot 3^b$ ($a, b \in \mathbb{N}_0$), where at least one of numbers a and b differs from 0. Let's introduce the following symbols.

1. (k, l) , $k, l \in \mathbb{N}_0$, $k^2 + l^2 \geq 1$ will denote number $2^k \cdot 3^l$. This k can be any nonnegative integer – it has no connection to k from the text of the problem.
2. $g: \mathbb{N}_0 \times \mathbb{N}_0 \rightarrow \mathbb{N}_0$, $g(k, l) = f(2^k \cdot 3^l)$, $k^2 + l^2 \geq 1$.

Now we can write that Paralampius can jump from pair (k, l) to pair

- $(k+1, l)$ (always),
- $(k, l+1)$ (always),
- $(2k, 2l)$ (if $4k \geq a+1$ and $4l \geq b+1$ with pair (a, b) being the goal).

Let pair (a, b) in further text always mark the number which is Paralampius' goal. Let's show that Paralampius can jump according to the third rule once, at most. Let's assume the opposite, that is, that Paralampius has jump from pair (k, l) to pair $(2k, 2l)$ and after a couple of jumps, again from pair (k_1, l_1) to pair $(2k_1, 2l_1)$. It is obvious that none of these rules decrease any element in a pair. On the contrary, every rule increases at least one number in a pair, while the third rule increases (doubles) both numbers. So, it is true that $k_1 \geq 2k \geq \frac{a+1}{2}$, so $2k_1 \geq a+1 > a$, and, as it has been mentioned previously, none of the rules will decrease numbers in a pair, so Paralampius will never be able to reach pair (a, b) . Contradiction! So, Paralampius will jump according to the third rule at most once. If he never jumps according to the third rule, he can jump from $(0, 0)$ to (a, b) in

$$\binom{a+b}{b}$$

ways. Further, if Paralampius jumps once according to the third rule, for instance from pair (k, l) to pair $(2k, 2l)$, then it must hold for k and l that $4k \geq a+1$ and $4l \geq b+1$ (condition of the problem), but also (to keep Paralampius from "overjumping" the goal, because, as we have shown, he can not return) $2k \leq a$ and $2l \leq b$. So, Paralampius can come from $(0, 0)$ to (a, b) , if he jumps once from (k, l) to $(2k, 2l)$ in

$$\binom{k+l}{l} \cdot \binom{a+b-2k-2l}{b-2l}$$

ways. According to that, Paralampius can come from $(0, 0)$ to (a, b) in, altogether,

$$g(a, b) = \binom{a+b}{b} + \sum_{\frac{a+1}{4} \leq k \leq \frac{a}{2}} \sum_{\frac{b+1}{4} \leq l \leq \frac{b}{2}} \binom{k+l}{l} \cdot \binom{a+b-2k-2l}{b-2l}$$

ways.

(2 points)

Now our problem boils down to showing that for each natural number n there exists a natural number m_0 such that for every natural number $2^a \cdot 3^b > m_0$ ($a, b \in \mathbb{N}_0$, $a^2 + b^2 \geq 1$), it holds that

$$g(a, b) < 2^{a+b-n}.$$

Let us now show three lemmas.

Lemma 1. For all natural numbers x and y , $\binom{x}{y} \leq 2^{x-1}$ holds.

Proof. We know that $\binom{x}{y} \leq \binom{x}{\lfloor \frac{x}{2} \rfloor}$, so it suffices to prove that, for every natural number x , $\binom{x}{\lfloor \frac{x}{2} \rfloor} \leq 2^{x-1}$. We will show this using mathematical induction. For $x = 1$, $\binom{1}{0} \leq 2^{1-1}$. Let's assume that for some $x \in \mathbb{N}$, $\binom{x}{\lfloor \frac{x}{2} \rfloor} \leq 2^{x-1}$. Then,

$$\binom{x+1}{\lfloor \frac{x+1}{2} \rfloor} = \binom{x}{\lfloor \frac{x+1}{2} \rfloor} + \binom{x}{\lfloor \frac{x-1}{2} \rfloor} \leq 2^{x-1} + 2^{x-1} = 2^x.$$

We shall notice that equality is only possible if $x = 1$ or $x = 2$. This proves our first lemma.

(1 point)

■

Lemma 2. There exists a natural number c_1 such that, if $a > c_1$ or $b > c_1$ (then $a + b > c_1$), then

$$(a+3)(b+3) < 2^{\frac{a+b}{4}}.$$

Proof. If we fix $a + b$, then the left side reaches its maximum for $a = b$. Let then be $M = a + b + 6$. Then,

$$M = (a+3) + (b+3) \geq [\text{using AM-GM inequality}] \geq 2\sqrt{(a+3)(b+3)}.$$

So, it suffices to show that there exists a natural number c_1 such that for every natural number $x > c_1$ it holds that

$$(x+3)^2 < 2^{\frac{x}{2}} \iff (x+3)^4 < 2^x.$$

It is obvious that the left side of this inequality “grows more slowly” than the right side, so it is enough to show that there exists at least one such natural number x . We can directly see that, for example, $x = 20$ has this property.

(1 point)

■

Lemma 3. For every natural number n there exists natural number c such that, if $a > c$ or $b > c$ (then $a + b > c$), then

$$\binom{a+b}{b} < 2^{a+b-n-1}.$$

Proof. We know that

$$\binom{a+b}{b} \leq \binom{a+b}{\lfloor \frac{a+b}{2} \rfloor},$$

so, in general,

$$\binom{a+b}{b} \leq \binom{2x}{x}, \quad \text{where } x = \left\lceil \frac{a+b}{2} \right\rceil.$$

Now, it is enough to show that for each natural number n there exists natural number c such that, if $x > c$, then

$$\binom{2x}{x} < 2^{2x-n-1}.$$

Further, let us notice that $\binom{2x+2}{x+1} < 4 \cdot \binom{2x}{x} \iff 2 > 0$, and $2^{2x+2-n-1} = 4 \cdot 2^{2x-n-1}$. So, it inductively follows that, if we show that there exists $c \in \mathbb{N}$ such that $\binom{2c}{c} < 2^{2c-n-1}$, then, for every natural number $x > c$, $\binom{2x}{x} < 2^{2x-n-1}$ will hold. Let us now prove that such $c \in \mathbb{N}$ exists.

$$\begin{aligned} & \binom{2c}{c} < 2^{2c-n-1} \\ \iff & \frac{(2c)!}{c! \cdot c!} < 2^{2c-n-1} \\ \iff & \frac{(2c)(2c-1)(2c-2) \cdots 2 \cdot 1}{c^2(c-1)^2(c-2)^2 \cdots 2^2 \cdot 1^2} < 2^{2c-n-1} \\ \iff & \frac{2^c \cdot c(c-1)(c-2) \cdots 2 \cdot 1 \cdot (2c-1)(2c-3) \cdots 3 \cdot 1}{c^2(c-1)^2 \cdots 2^2 \cdot 1^2} < 2^{2c-n-1} \\ \iff & \frac{(2c-1)(2c-3) \cdots 3 \cdot 1}{c(c-1) \cdots 2 \cdot 1} < 2^{c-n-1} \\ \iff & \frac{(2c-1)(2c-3) \cdots 3 \cdot 1}{2^c \cdot c(c-1) \cdots 2 \cdot 1} < 2^{-n-1} \\ \iff & \frac{(2c)(2c-2)(2c-4) \cdots 4 \cdot 2}{(2c-1)(2c-3) \cdots 3 \cdot 1} > 2^{n+1} \\ \iff & \left(1 + \frac{1}{2c-1}\right) \left(1 + \frac{1}{2c-3}\right) \cdots \left(1 + \frac{1}{1}\right) > 2^{n+1}. \end{aligned}$$

Obviously $\left(1 + \frac{1}{2c-1}\right) \left(1 + \frac{1}{2c-3}\right) \cdots \left(1 + \frac{1}{1}\right) \geq \frac{1}{2c-1} + \frac{1}{2c-3} + \cdots + \frac{1}{1}$ for every natural number c , so it suffices to show that there exists a natural number c such that

$$\frac{1}{2c-1} + \frac{1}{2c-3} + \cdots + \frac{1}{1} > 2^{n+1}.$$

Since $\frac{1}{2c-1} + \frac{1}{2c-3} + \cdots + \frac{1}{1} \geq \frac{1}{2c} + \frac{1}{2c-2} + \cdots + \frac{1}{2}$, that is,

$$\frac{1}{2c-1} + \frac{1}{2c-3} + \cdots + \frac{1}{1} \geq \frac{\frac{1}{c} + \frac{1}{c-1} + \cdots + \frac{1}{1}}{2},$$

it is enough for us to show that there exists a natural number c such that

$$\frac{1}{c} + \frac{1}{c-1} + \cdots + \frac{1}{1} > 2^{n+2}.$$

Let's notice that for every nonnegative integer t , $\frac{1}{2^t+1} + \frac{1}{2^t+2} + \cdots + \frac{1}{2^{t+1}} \geq \frac{1}{2}$. This follows from the fact that, on the left side of the inequality, we have 2^t summands, each one of which is larger or equal than $\frac{1}{2^{t+1}}$. Now it directly follows that

$$\frac{1}{2^{2n+3}} + \frac{1}{2^{2n+3}-1} + \frac{1}{2^{2n+3}-2} + \cdots + \frac{1}{1} > 2^{n+2}.$$

With this, we have finally proven this lemma. (3 points) ■

Now, we wish to show that for every natural number n there exists a natural number t such that, if $a > t$ or $b > t$, then

$$\sum_{\frac{a+1}{4} \leq k \leq \frac{a}{2}} \sum_{\frac{b+1}{4} \leq l \leq \frac{b}{2}} \binom{k+l}{l} \cdot \binom{a+b-2k-2l}{b-2l} < 2^{a+b-n-1}.$$

Further, let's fix n . Let t be such natural number that $t > c_1$ and $t > c$, where c_1 and c are numbers of **Lemma 2**, that is, from **Lemma 3**. Let us observe numbers defined by pair (a, b) , where $a + 1 > 4t$ or $b + 1 > 4t$. Then, if Paralampius jumps on his travel from pair (k, l) to pair $(2k, 2l)$, for numbers k and l it holds that $k + l > t$ because $k \geq \frac{a+1}{4}$ and $l \geq \frac{b+1}{4}$, so $k > t$ or $l > t$. So, because of

Lemma 3, $\binom{k+l}{l} < 2^{k+l-n-1}$. According to **Lemma 1**, we have that for all $x, y \in \mathbb{N}$, it holds that $\binom{x}{y} \leq 2^{x-1}$, so, because of that,

$$\sum_{\frac{a+1}{4} \leq k \leq \frac{a}{2}} \sum_{\frac{b+1}{4} \leq l \leq \frac{b}{2}} \binom{k+l}{l} \cdot \binom{a+b-2k-2l}{b-2l} < \sum_{\frac{a+1}{4} \leq k \leq \frac{a}{2}} \sum_{\frac{b+1}{4} \leq l \leq \frac{b}{2}} 2^{k+l-n-1} \cdot 2^{a+b-2k-2l-1}.$$

Since $k > \frac{a}{4}$ and $l > \frac{b}{4}$, we can see that it suffices to show that (we set k to be exactly equal to $\frac{a}{4}$ and the same for l)

$$\sum_{\frac{a+1}{4} \leq k \leq \frac{a}{2}} \sum_{\frac{b+1}{4} \leq l \leq \frac{b}{2}} 2^{\frac{3}{4}a + \frac{3}{4}b - n - 2} < 2^{a+b-n-1}.$$

It is obvious that there exists at most $\frac{a}{2} - \frac{a+1}{4} + 1 = \frac{a+3}{4}$ possible numbers k . The same is true for numbers l . So, it is enough for us to show

$$(a+3)(b+3) \cdot 2^{\frac{3}{4}a + \frac{3}{4}b - n - 5} < 2^{a+b-n}.$$

Since $a > 4t - 1$ or $b > 4t - 1$, then $a > t$ or $b > t$ and, since $t > c_1$, where c_1 is the number from **Lemma 2**, it is obvious that $a > c_1$ or $b > c_1$, so, because of **Lemma 2**, wanted inequality directly follows. With this we have shown that for every natural number n , there exists a natural number t such that, if $a > t$ or $b > t$, then

$$\sum_{\frac{a+1}{4} \leq k \leq \frac{a}{2}} \sum_{\frac{b+1}{4} \leq l \leq \frac{b}{2}} \binom{k+l}{l} \cdot \binom{a+b-2k-2l}{b-2l} < 2^{a+b-n-1}.$$

In our case, this natural number t is such that $t > 4c_1$ and $t > 4c$, where c_1 and c are numbers from **Lemma 2**, that is, from **Lemma 3**. (2 points)

It is obvious that, for such natural number t , because of **Lemma 3**, the following holds:

$$\binom{a+b}{b} < 2^{a+b-n-1}.$$

By adding the last two inequalities (we have shown that they hold), we reach that

$$g(a, b) < 2^{a+b-n},$$

what we have wanted to show. That is, we have shown that for every natural number n there exists a natural number m_0 such that for every natural number $2^a \cdot 3^b > m_0$ ($a, b \in \mathbb{N}_0$, $a^2 + b^2 \geq 1$), it holds that


$$g(a, b) < 2^{a+b-n}.$$



Problem 3. A convex n -gon ($n \in \mathbb{N}$, $n > 2$) is given in the plane. Its area is less than 1. For each point X of this plane, we shall denote with $F(X)$ the area of the convex hull of point X and a given n -gon (the area of the minimal convex polygon which includes both the point X and a given n -gon). Prove that the set of points for which $F(X) = 1$ is a convex polygon with $2n$ sides or less.

Solution. Let S be the set of all points T of the given plane such that $F(T) = 1$. It is obvious that all these points T are outside the given n -gon. Area of the given n -gon is less than 1, and when point T belongs to the interior (or the edge) of the given n -gon, it itself becomes our wanted convex hull. (1 point)

Point T is outside the n -gon if there exists a line which passes through T such that it has no common points with the n -gon. Let's observe the following image.



Let A_0, A_1, \dots, A_{n-1} be vertices of the given n -gon. Also, let $A_j = A_{(j \bmod n)}$ ($j \in \mathbb{Z}$), where $(j \bmod n) = x$ is a number from the set $\{0, 1, \dots, n-1\}$ such that $j \equiv x \pmod{n}$. We say that a diagonal of the given n -gon is *visible* if the entire n -gon is inside the angle with its vertex in T whose chord is that diagonal. For example, (observe image) diagonal $\overline{A_i A_k}$ ($i, k \in \mathbb{Z}$), with respect to X , is visible. (1 point)

$\overline{A_i A_k}$ is a visible diagonal. Inside the triangle $A_i X A_k$ there are vertices $A_{i+1}, A_{i+2}, \dots, A_{k-1}$ and outside of it all the remaining vertices of the given n -gon. So, the area $F(X)$ is equal to the sum of areas of the triangle $A_i X A_k$ and polygon $A_k A_{k+1} \dots A_{i-1} A_i$. (1 point)

Let l be a line through X parallel to the line $A_i A_k$ and let A and B be the intersections of l and lines $A_{i-1} A_i$ and $A_{k+1} A_k$, respectively. (1 point)

If $F(X) = 1$, then X is an element of S . Then, it is obvious that all points T that are on line l and for which $\overline{A_i A_k}$ is a visible diagonal are also elements of set S . Furthermore, diagonal $\overline{A_i A_k}$ is no longer visible to the point T as T moves on l and becomes colinear with some side of the given n -gon. So, all points T of line l that are possibly in S (which they are if X is in S) are on the segment \overline{AB} . (2 points)

Now we can easily deduce the following conclusion: By observing all points $T \in S$ we can see that, as they “go around” the given n -gon, with them also rotate their respective visible diagonals. Every point of the given n -gon can, therefore, become one endpoint of some of the visible diagonals (visible diagonals interesting to us – those who are visible with respect to some point from the set S)

exactly once, and only exactly once stop being the endpoint. Because of that, there can be at most $2n$ interesting visible diagonals. It is clear that we will get a convex polygon – as point T rotates around the given n -gon, segments \overline{AB} rotate with her and one always “connects” itself to the other one.

(5 points) ■

Problem 4. Prove that for every natural number k , there exist infinitely many natural numbers n such that

$$\frac{n - d(n^r)}{r} \in \mathbb{Z}, \text{ for every } r \in \{1, 2, \dots, k\}.$$

Here, $d(x)$ denotes the number of natural divisors of a natural number x , including 1 and x itself.
(Melkior Ornik)

Solution. Let us first prove the following lemma.

Lemma 1. $d(n^r) \equiv 1 \pmod{r}$.

Proof. Let $n = p_1^{\alpha_1} \cdot p_2^{\alpha_2} \cdot \dots \cdot p_i^{\alpha_i}$, where $p_1 < p_2 < \dots < p_i$ are all prime divisors of the natural number n and $i, \alpha_1, \alpha_2, \dots, \alpha_i$ are natural numbers, be the prime factorization of n . Then,

$$n^r = p_1^{r\alpha_1} \cdot p_2^{r\alpha_2} \cdot \dots \cdot p_i^{r\alpha_i},$$

that is, $d(n^r) = (r\alpha_1 + 1) \cdot (r\alpha_2 + 1) \cdot \dots \cdot (r\alpha_i + 1)$. (1 point)

Clearly, for every $j \in \{1, 2, \dots, i\}$, the following holds: $r\alpha_j + 1 \equiv 1 \pmod{r}$, so

$$d(n^r) \equiv (r\alpha_1 + 1) \cdot (r\alpha_2 + 1) \cdot \dots \cdot (r\alpha_i + 1) \equiv 1^i \equiv 1 \pmod{r}.$$

This has proven the lemma. (1 point) ■

Now, our problem (because of **Lemma 1**) boils down to showing that for every natural number k there exists infinitely many natural numbers n such that

$$r \mid n - 1, \quad \forall r \in \{1, 2, \dots, k\}.$$

Let $n = m \cdot k! + 1$, m being a natural number. (1 point)

Now it is obvious that $n \equiv m \cdot k! + 1 \equiv [\text{because of } r \mid k!] \equiv 1 \pmod{r}$ for every $r \in \{1, 2, \dots, k\}$.

Because of that, for this choice of number n we have $r \mid n - 1$ for every $r \in \{1, 2, \dots, k\}$. (1 point)

As m can be any natural number, for every natural number k there exists infinitely many natural numbers n such that $r \mid n - 1$ for every $r \in \{1, 2, \dots, k\}$. This is, because of **Lemma 1**, equivalent to the problem claim. (6 points) ■