

Yugoslav Team Selection Tests 2002

Selection Test for Balkan MO

Bečići, April 21

1. A man standing at the point $(1, 1)$ in the coordinate plane wants to find an object that lies at some point (α, β) , where $\alpha \in \{1, 2, \dots, m\}$ and $\beta \in \{1, 2, \dots, n\}$. After finding the object, he will return to the starting point. Find the minimum time that is always sufficient for doing this job in terms of m and n . We assume that the man does not know at which point the object lies, and that he can move in any direction with velocity not exceeding 1.
2. Let p be the semiperimeter of a triangle ABC . Points E and F are taken on line AB such that $CE = CF = p$. Show that the circumcircle of the triangle EFC is tangent to the excircle of the triangle ABC that meets the side AB .
3. The sequence $(x_n)_{n=2}^{\infty}$ is defined by $x_2 = x_3 = 1$ and for $n \geq 3$,

$$(n+1)(n-2)x_{n+1} = n(n^2 - n - 1)x_n - (n-1)^3x_{n-1}.$$

Prove that x_n is an integer if and only if n is a prime number.

Selection Test for IMO

Belgrade, June ?

1. Let a, b, c be nonnegative numbers such that $a^2 + b^2 + c^2 + abc = 4$. What is the maximum value of $a + b + c + abc$?
2. Let $ABCD$ be a convex quadrilateral with $\angle DAB = \angle ABC = \angle BCD$. Let O and H be the circumcenter and the orthocenter of triangle ABC , respectively. Prove that the points O, H , and D are collinear.
3. For any positive integer n , we denote by $f(n)$ the number of possible choices for plus and minus signs in the expression $1 \pm 2 \pm \dots \pm n$ such that the value of the expression is 0.
 - (a) Prove that $f(n) = 0$ for $n \equiv 1, 2 \pmod{4}$.
 - (b) Prove that $f(n) \geq 2^{\frac{n}{2}-1}$ for $n \equiv 0, 3 \pmod{4}$.